

<b><i>Eucalyptus kruseana</i> (Book-Leaf Mallee, Kruse's Mallee) -- FLORIDA</b>		<b>Answer</b>	<b>Score</b>
1.01	Is the species highly domesticated?	n	0
1.02	Has the species become naturalised where grown?		
1.03	Does the species have weedy races?		
2.01	Species suited to FL climates (USDA hardiness zones; 0-low, 1-intermediate, 2-high)	2	
2.02	Quality of climate match data (0-low; 1-intermediate; 2-high)	2	
2.03	Broad climate suitability (environmental versatility)	y	1
2.04	Native or naturalized in regions with an average of 11-60 inches of annual precipitation	n	0
2.05	Does the species have a history of repeated introductions outside its natural range?	n	
3.01	Naturalized beyond native range	n	0
3.02	Garden/amenity/disturbance weed	n	0
3.03	Weed of agriculture	n	0
3.04	Environmental weed	n	0
3.05	Congeneric weed	y	2
4.01	Produces spines, thorns or burrs	n	0
4.02	Allelopathic	?	
4.03	Parasitic	n	0
4.04	Unpalatable to grazing animals	?	
4.05	Toxic to animals	?	
4.06	Host for recognised pests and pathogens		
4.07	Causes allergies or is otherwise toxic to humans		
4.08	Creates a fire hazard in natural ecosystems	?	
4.09	Is a shade tolerant plant at some stage of its life cycle	?	
4.10	Grows on infertile soils (oligotrophic, limerock, or excessively draining soils). North & Central Zones: infertile soils; South Zone: shallow limerock or Histisols.	y	1
4.11	Climbing or smothering growth habit	n	0
4.12	Forms dense thickets	n	0
5.01	Aquatic	n	0
5.02	Grass	n	0
5.03	Nitrogen fixing woody plant	n	0
5.04	Geophyte	n	0
6.01	Evidence of substantial reproductive failure in native habitat		
6.02	Produces viable seed	y	1
6.03	Hybridizes naturally	y	1
6.04	Self-compatible or apomictic		
6.05	Requires specialist pollinators	n	0
6.06	Reproduction by vegetative propagation		
6.07	Minimum generative time (years)	3	0

7.01	Propagules likely to be dispersed unintentionally (plants growing in heavily trafficked areas)		
7.02	Propagules dispersed intentionally by people	y	1
7.03	Propagules likely to disperse as a produce contaminant		
7.04	Propagules adapted to wind dispersal	n	-1
7.05	Propagules water dispersed	?	
7.06	Propagules bird dispersed	n	-1
7.07	Propagules dispersed by other animals (externally)	n	-1
7.08	Propagules dispersed by other animals (internally)	n	-1
8.01	Prolific seed production		
8.02	Evidence that a persistent propagule bank is formed (>1 yr)	n	-1
8.03	Well controlled by herbicides	?	
8.04	Tolerates, or benefits from, mutilation or cultivation	y	1
8.05	Effective natural enemies present in U.S.		
	<b>Total Score</b>		<b>3</b>
	<b>Implemented Pacific Second Screening</b>		<b>Yes</b>
	<b>Risk Assessment Results</b>		<b>Accept</b>

	Reference	Source data
1.01		Cultivated but no evidence of selection for reduced weediness.
1.02		Skip to 2.01
1.03		Skip to 2.01
2.01	1. PERAL NAPPFast Global Plant Hardiness ( <a href="http://www.nappfast.org/Plant_hardiness/NAPPFast%20Global%20zones/10-year%20climate/PLANT_HARDINESS_10YR%20lgnd.tif">http://www.nappfast.org/Plant_hardiness/NAPPFast%20Global%20zones/10-year%20climate/PLANT_HARDINESS_10YR%20lgnd.tif</a> ) & USDA Plant Hardiness Zone Map, 2012. Agricultural Research Service, U.S. Department of Agriculture. Accessed from <a href="http://planthardiness.ars.usda.gov">http://planthardiness.ars.usda.gov</a> . 2. USDA/ARS-GRIN [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland ( <a href="http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?15948">http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?15948</a> [Accessed: 8 May 2012]). 3. " <i>Eucalyptus kruseana</i> ." hortipedia.com. Hortipedia, 2011. Web. 21 May 2012. 4. Australian Native Plant Society (Australia) (ANPSA). <a href="http://anpsa.org.au/index.html">http://anpsa.org.au/index.html</a> . Accessed: 23 May 2012.	<b>No computer analysis was performed.</b> 1. Global plant hardiness zones 9-10; equivalent to USDA Hardiness zones 8b-10a (north, central, and south zones of Florida). 2. Native distribution: Westernn Australia and possibly in Queensland, Australia. 3. Hardy range: 9a-10a; native to Western Australia. 4. Restricted to three hilly areas of east Kalgoorlie, Western Australia.
2.02		<b>No computer analysis was performed.</b> 1. Native range is well known; refer to 2.01 source data.
2.03	1. Köppen-Geiger climate map ( <a href="http://www.hydrol-earth-syst-sci.net/11/1633/2007/hess-11-1633-2007.pdf">http://www.hydrol-earth-syst-sci.net/11/1633/2007/hess-11-1633-2007.pdf</a> ).	1. Native distribution appears to be in at least three climatic groups (BSk, Csa, Cfa).
2.04	1. Australia's Virtual Herbarium. 2009. <a href="http://chah.gov.au/avh/index.jsp">http://chah.gov.au/avh/index.jsp</a> . Accessed: 9 May 2012.	1. 200 mm-800 mm (7.9"-31.5").
2.05		No evidence.
3.01		No evidence.
3.02		No evidence.
3.03		No evidence.
3.04		No evidence.
3.05	1. Holm, L. et al. <i>A Geographical Atlas of World Weeds</i> . John Wiley and Sons, New York. 1979.	1. The following <i>eucalypts</i> are considered principal weeds in Australia (principal weed in this context is ranked according to the importance of the weed and is usually referring to about the five most troublesome species for the crop): <i>E. cambageana</i> , <i>E. ferruginea</i> , <i>E. gracilis</i> , <i>E. marginata</i> , <i>E. miniata</i> , <i>E. pilularis</i> , <i>E. populnea</i> , <i>E. tetradonta</i> .
4.01		No evidence.

4.02	<p>1. Anonymous. 2009. "Focus on Eucalypts." SAPIA NEWS No. 12. ARC-Plant Protection Research Institute, South Africa. 2. Anonymous. October 2010. Scotland, Forestry Commission. Interim Guidance on the Grant Aiding and Planting of Eucalypts in Scotland. Accessed: 1 June 2012. 3. Rejmánek, M. &amp; D.M. Richardson. 2011. Eucalypts (203-209). In D. Simberloff &amp; M. Rejmánek, eds. <i>Encyclopedia of Biological Invasions</i>. Berkeley: University of California Press.</p>	<p>1. It is likely that most Eucalypts are allelopathic-having the potential to suppress understory plants through chemical inhibitors that leach into the soil. 2. There are many reports in global literature of toxic inhibition of germination and growth of other plant species (allelopathic effects), which inhibits the growth of an understory. 3. Concerns expressed about suppression of ground vegetation due to possible allelopathic effects. Allelopathic effects are widely reported and these reports are largely based on laboratory bioassays. If not chemical inhibition then at least accumulation of dead material of the floor of eucalypt plantations hinders regeneration of native species.</p>
4.03		No evidence.
4.04	<p>1. United States Department of Agriculture Permit applications 08-11-106rm and 08-014-101rm received from ArborGen LLC. Field testing of genetically engineered <i>E. grandis</i> X <i>E. urophylla</i> (<a href="http://www.aphis.usda.gov/brs/aphisdocs/08_014101rm_ea2.pdf">http://www.aphis.usda.gov/brs/aphisdocs/08_014101rm_ea2.pdf</a> [Accessed: 8/19/2010]).</p>	<p>1. Eucalyptus species are known to produce chemical compounds that are required by the plant for defense against herbivores and pathogens.</p>
4.05	<p>1. <i>Medicinal Plants for Livestock: Eucalyptus spp.</i> Cornell University, Department of Animal Science. <a href="http://www.ansci.cornell.edu/plants/medicinal/eucalyp.html">http://www.ansci.cornell.edu/plants/medicinal/eucalyp.html</a>. 1 June 2012.</p>	<p>1. "<i>Eucalyptus spp.</i> contain high levels of phenolics and terpenoids which can be toxic. Animals such as the koala which eat Eucalyptus have developed methods for detoxifying the compounds in the liver. In addition, they have bacteria that degrade tannin-protein complexes. Most animals do not have this ability."</p>
4.06		
4.07		
4.08	<p>1. Gill, A.M. "Eucalypts and fires: interdependent or independent?" In: <i>Eucalypt ecology: individuals to ecosystems</i>. Ed. J.E. Williams &amp; J. Woinarski. Cambridge, New York: Cambridge University Press, 1997. 2. Rejmánek, M. &amp; D.M. Richardson. 2011. Eucalypts (203-209). In D. Simberloff &amp; M. Rejmánek, eds. <i>Encyclopedia of Biological Invasions</i>. Berkeley: University of California Press.</p>	<p>1. Eucalypts often are the major source of fuel for fires, but not always. 2. Leaves of eucalypts are relatively slow to breakdown and have a high volatile oil content, which contributes to the severity of fire events in their native Australia.</p>
4.09	<p>1. Rejmánek, M. &amp; D.M. Richardson. 2011. Eucalypts (203-209). In D. Simberloff &amp; M. Rejmánek, eds. <i>Encyclopedia of Biological Invasions</i>. Berkeley: University of California Press.</p>	<p>1. Shade-tolerant sub-canopy [<i>Eucalyptus</i>] species are not known.</p>

4.10	1. " <i>Eucalyptus kruseana</i> ." hortycopia.com. Hortycopia, 2011. Web. 30 May 2012. 2. <i>Eucalyptus kruseana</i> . FloraBase: Flora of Western Australia. Accessed 1 June 2012. <a href="http://florabase.dec.wa.gov.au/browse/profile/5687">http://florabase.dec.wa.gov.au/browse/profile/5687</a>	1. "Suitable soil is well-drained/loamy or sandy. The pH preference is a neutral soil." 2. "Sandy loam. Granite outcrops & hills"
4.11	1. " <i>Eucalyptus kruseana</i> ." hortycopia.com . Hortycopia, 2011. Web. 21 May 2012.	1. "Tree, shrub, woody plant, up to 3 m (10')".
4.12	1. " <i>Eucalyptus kruseana</i> ." hortycopia.com . Hortycopia, 2011. Web. 21 May 2012.	1. Up to 3 m (10') tall.
5.01	1. Australian Native Plant Society (Australia) (ANPSA). <a href="http://anpsa.org.au/index.html">http://anpsa.org.au/index.html</a> . Accessed: 23 May 2012.	1. Occurs on rocky hills in an arid environment.
5.02	1. USDA/ARS-GRIN [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland ( <a href="http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?15948">http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?15948</a> [Accessed: 8 May 2012]).	1. Family: <i>Myrtaceae</i> .
5.03	1. USDA/ARS-GRIN [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland ( <a href="http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?15948">http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?15948</a> [Accessed: 8 May 2012]).	1. Family: <i>Myrtaceae</i> .
5.04	1. " <i>Eucalyptus kruseana</i> ." hortycopia.com. Hortycopia, 2011. Web. 21 May 2012. 2. Pacific Island Ecosystems at Risk (PIER). <a href="http://www.hear.org">http://www.hear.org</a> . Via: <a href="http://plantsdatabase.com/go/55096/">http://plantsdatabase.com/go/55096/</a> .	1. "Tree, shrub, woody plant".
6.01		
6.02	1. Australian Native Plant Society (Australia) (ANPSA). <a href="http://anpsa.org.au/index.html">http://anpsa.org.au/index.html</a> . Accessed: 23 May 2012. 2. Pacific Island Ecosystems at Risk (PIER). <a href="http://www.hear.org">http://www.hear.org</a> . Via: Chippendale, George McCartney. <i>Eucalypts of the Western Australian goldfields : (and the adjacent wheatbelt)</i> Australian Government Publishing Service for the Minister for Primary Industry, Canberra 1973. 218 pp. p.84	1. Propagation is from seed which germinates readily. 2. The average of two samples of seed was 134 viable seeds per gram of fertile and sterile seeds.

6.03	<p>1. Pacific Island Ecosystems at Risk (PIER). <a href="http://www.hear.org">http://www.hear.org</a>. Via: Grayling, P.M. &amp; M.I.H Brooker. 1996. Evidence for the identity of the hybrid <i>Eucalyptus 'brachyphylla'</i> (<i>Myrtaceae</i>) from morphology and essential-oil composition. <i>Australian Journal of Botany</i>, 44(1): 1-13.</p> <p>2. Slee, A.V, M.I.H. Brooker, S.M. Duffy, J.G. West. 2006. Centre for Plant Biodiversity Research. EUCLID Eucalypts of Australia, 3rd Edition. <a href="http://www.anbg.gov.au/cpbr/cd-keys/Euclid/sample/html/index.htm">http://www.anbg.gov.au/cpbr/cd-keys/Euclid/sample/html/index.htm</a>. Accessed: 23 May 2012.</p>	<p>1. The putative interspecific hybrid known as <i>Eucalyptus brachyphylla</i>, which is known to occur at two sites near Kalgoorlie, Western Australia, contains the non-terpenoid ester 4-methyl-2-pentyl acetate and the related alcohol 4-methylpentan-2-ol, in its leaves. These compounds are otherwise known to occur in the leaves of only four <i>Eucalyptus</i> species and several putative interspecific hybrids. This, with the results of multivariate analyses of morphometric and oil composition data, adds considerable weight to earlier suggestions that <i>E. 'brachyphylla'</i> is a hybrid of <i>E. loxophleba</i> and <i>E. kruseana</i>. 2. There are a few well known formally named hybrids where both parents have been identified and are seen in the field, e.g., <i>E. brachyphylla</i> (<i>E. kruseana</i> X <i>E. loxophleba</i> subsp. <i>lissophloia</i>).</p>
6.04		
6.05	<p>1. "<i>Eucalyptus kruseana</i> ." horticoxia.com. Horticoxia, 2011. Web. 21 May 2012. 2. Pacific Island Ecosystems at Risk (PIER). <a href="http://www.hear.org">http://www.hear.org</a>. Via: <a href="http://plantsdatabase.com/go/55096/">http://plantsdatabase.com/go/55096/</a>.</p>	<p>1. Attracts birds and butterflies. 2. This plant is attractive to bees, butterflies and/or birds.</p>
6.06		
6.07	<p>1. Pacific Island Ecosystems at Risk (PIER). <a href="http://www.hear.org">http://www.hear.org</a>. Via: (1)<a href="http://www.australiaplants.com/Eucalyptus_kruseana.htm">http://www.australiaplants.com/Eucalyptus_kruseana.htm</a>.</p>	<p>1. usually starts flowering in the third year.</p>
7.01		
7.02		<p>Species is being considered for introduction as a biomass crop.</p>
7.03		

7.04	<p>1. Pacific Island Ecosystems at Risk (PIER). <a href="http://www.hear.org">http://www.hear.org</a>. Via: Chippendale, George McCartney. Eucalypts of the Western Australian goldfields : (and the adjacent wheatbelt) Australian Government Publishing Service for the Minister for Primary Industry, Canberra 1973. 218 pp. p.84. 2. Potts, B. 1990. The response of eucalypt populations to a changing environment. Tasforests, December: 179-193. 3. Cremer, K.W. 1977. Distance of seed dispersal in Eucalypts estimated from seed weights. Australian Forest Research, 7(4): 225-228. 4. Rejmánek, M. &amp; D.M. Richardson. 2011. Eucalypts (203-209). In: D. Simberloff &amp; M. Rejmánek, eds. Encyclopedia of Biological Invasions. Berkeley: University of California Press.</p>	<p>No adaptations for wind dispersal (i.e., lacks wings). 1. The fertile seeds are brown, oblong, crescent-shaped, 1-2mm long and with a net-like pattern. The sterile seeds are red-brown, wedge-shaped, about 1mm long and with a faint, net-like pattern. 2. Seed dispersal in most eucalypt species is mainly by wind and gravity. 3. Wind is probably the only important agent of seed dispersal in the eucalypts, except possibly in species growing on river margins or flood plains where water could also transport the seed. 4. Relatively limited seed dispersal; planted eucalypts are very small and have no adaptations for dispersal (wings or fleshy). The passive release of seeds is undoubtedly aided by wind; however all rigorous studies of eucalypt seed dispersal and seedling spatial distribution show that in general seeds are dispersed over quite short distances that are in agreement with measurement of terminal descent velocity.</p>
7.05	<p>1. Rejmánek, M. &amp; D.M. Richardson. 2011. Eucalypts (203-209). In D. Simberloff &amp; M. Rejmánek, eds. <i>Encyclopedia of Biological Invasions</i> . Berkeley: University of California Press.</p>	<p>1. Eucalypts should not be planted near rivers/streams. Temporarily flooded or eroded river/stream banks are suitable habitat for spontaneous establishment of seedlings. Additionally, their seeds can be dispersed for long distances by running water.</p>
7.06	<p>1. Southern, S.G. et al. 2004. Review of gene movement by bats and birds and its potential significance for eucalypt plantation forestry. <i>Australian Forestry</i> , 67(1): 44-53.</p>	<p>1. Dispersal in animal droppings does not occur, although many birds eat eucalypt seed, because the seed does not survive passage through the alimentary canal of mammals and birds (Joseph 1986).</p>
7.07	<p>1. Pacific Island Ecosystems at Risk (PIER). <a href="http://www.hear.org">http://www.hear.org</a>. Via: Chippendale, George McCartney. Eucalypts of the Western Australian goldfields : (and the adjacent wheatbelt) Australian Government Publishing Service for the Minister for Primary Industry, Canberra 1973. 218 pp. p.84</p>	<p>No adaptations that would suggest that it could attach itself externally to animals. 1. The fertile seeds are brown, oblong, crescent-shaped, 1-2mm long and with a net-like pattern. The sterile seeds are red-brown, wedge-shaped, about 1mm long and with a faint, net-like pattern. The average of two samples of seed was 134 viable seeds per gram of fertile and sterile seeds.</p>
7.08	<p>1. Southern, S.G. et al. 2004. Review of gene movement by bats and birds and its potential significance for eucalypt plantation forestry. <i>Australian Forestry</i> , 67(1): 44-53.</p>	<p>1. Dispersal in animal droppings does not occur, although many birds eat eucalypt seed, because the seed does not survive passage through the alimentary canal of mammals and birds (Joseph 1986).</p>
8.01		

8.02	1. Rejmánek, M. & D.M. Richardson. 2011. Eucalypts (203-209). In D. Simberloff & M. Rejmánek, eds. <i>Encyclopedia of Biological Invasions</i> . Berkeley: University of California Press.	1. Eucalypt seeds do not have dormancy and seed storage in the soil lasts less than a year.
8.03	1. Rejmánek, M. & D.M. Richardson. 2011. Eucalypts (203-209). In : D. Simberloff & M. Rejmánek, eds. <i>Encyclopedia of Biological Invasions</i> . Berkeley: University of California Press.	1. Triclopyr or glyphosate applied to freshly cut stumps can greatly reduce resprouting.
8.04	1. Australian Native Plant Society (Australia) (ANPSA). <a href="http://anpsa.org.au/index.html">http://anpsa.org.au/index.html</a> . Accessed: 23 May 2012. 2. Pacific Island Ecosystems at Risk (PIER). <a href="http://www.hear.org">http://www.hear.org</a> . Via: (a) Chippendale, George McCartney. Eucalypts of the Western Australian goldfields : (and the adjacent wheatbelt) Australian Government Publishing Service for the Minister for Primary Industry, Canberra 1973. 218 pp. p.84 (b) <a href="http://www.australiaplants.com/Eucalyptus_kruseana.htm">http://www.australiaplants.com/Eucalyptus_kruseana.htm</a> ; <a href="http://www.australiaplants.com/Eucalyptus_Info.htm">http://www.australiaplants.com/Eucalyptus_Info.htm</a>	1. It forms a lignotuber and would be expected to respond to pruning. 2. (a) "Light pruning only is recommended, as some cultivated specimens have not grown well after heavy pruning." (b) "lignotuber" a modified root system and enables the plant to produce new shoots even after it has been destroyed above ground (generally by fire or cold)
8.05		